

	Label Name	Generic Name	Prior-Use Storage	In-Use Storage	When to Inject	Duration	Pen Priming & Max Injectable Dose
<b>Rapid Acting</b>	Apidra U-100 Vial	I. glulisine	Refrig until expiration date or RT for up to 28 days	Refrig or RT for up to 28 days	Within 15 mins AC or within 20 min after starting a meal	3 to 4 hours	-
	<a href="#">Apidra U-100 SoloStar</a>			RT for up to 28 days Do not refrigerate pens or refill cartridges			Prime: 2 units Max: 80 units
	Novolog U-100 Vial	I. aspart		Refrig or RT for up to 28 days	Within 5-10 mins AC	3 to 7.2 hours	-
	<a href="#">Novolog U-100 FlexPen</a>			RT for up to 28 days Do not refrigerate			Prime: 2 units Max: 60 units
	Fiasp U-100 Vial	I. aspart		Refrig or RT for up to 28 days	At the start of a meal or within 20 mins after starting a meal	~5 to 7 hours	-
	<a href="#">Fiasp U-100 FlexTouch</a>						Prime: 2 units Max: 80 units
	Merilog U-100 Vial	I. aspart-szjj		Refrig or RT for up to 28 days	5-10 mins AC	3 to 5 hours	-
	<a href="#">Merilog U-100 Solostar</a>						RT for up to 28 days Do not refrigerate
	Admelog U-100 Vial	I. lispro		Refrig or RT for up to 28 days	Within 15 mins AC or immediately PC	~6.9 hours	-
	<a href="#">Admelog U-100 SoloStar</a>						RT for up to 28 days Do not refrigerate
	Kristy U-100 Vial	I. aspart-xjhz		Refrig or RT for up to 28 days	5-10 mins AC	3 to 5 hours	-
	<a href="#">Kirsty U-100 Pen</a>						RT for up to 28 days Do Not Refrigerate
	Humalog/Insulin Lispro U-100 Vial	I. lispro		Refrig or RT for up to 28 days	Within 15 mins AC or immediately PC	5.7 to 6.7 hours	-

All information obtained from Lexidrug or manufacturer package insert. Pen administration is illustrated at the bottom of each hyperlinked package insert.

RT: room temperature [68°F to 77°F (20°C to 25°C)]; Refrig: refrigeration [36°F - 46°F (2°C to 8°C)]. AC: before food; PC: after food.

	Label Name	Generic Name	Prior-Use Storage	In-Use Storage	When to Inject	Duration	Pen Priming & Max Injectable Dose
<b>Rapid Acting</b>	<a href="#">Humalog/Insulin Lispro U-100 KwikPen</a>	l. lispro	Refrig until expiration date or RT for up to 28 days	RT for up to 28 days Do not refrigerate pens or refill cartridges	Within 15 mins AC or immediately PC	5.7 to 6.7 hours	Prime: 2 units Max (U-100): 60 units Max (U-200): 60 units
	<a href="#">Humalog U-200 KwikPen</a>						
	Lyumjev U-100 Vial	l. lispro-aabc		Refrig or RT for up to 28 days	At start of meal or within 20 mins after starting meal	4.6 to 7.3 hours	-
	<a href="#">Lyumjev U-100 KwikPen</a>			RT for up to 28 days Do not refrigerate pens or refill cartridges			Prime: 2 units Max: 60 units
	<a href="#">Lyumjev U-200 KwikPen</a>						
<b>Short Acting</b>	Novolin-R U-100 Vial	l. regular	Refrig until expiration date or RT for up to 42 days	RT for up to 42 days Do not refrigerate	30 min AC	4 to 12 hours	-
	<a href="#">Novolin-R U-100 FlexPen</a>		Refrig until expiration date or RT for up to 28 days	RT for up to 28 days Do not refrigerate			Prime: 2 units Max: 60 units
	<a href="#">Humulin-R U-100 Vial</a>		Refrig until expiration date or RT for up to 31 days	RT or Refrig for up to 31 days			-
<b>Intermediate Acting</b>	Novolin-N U-100 Vial	l. NPH	Refrig until expiration date or RT for up to 42 days	RT for up to 42 days Do not refrigerate	Per written order	14 to 24 hours	-
	<a href="#">Novolin-N U-100 FlexPen</a>		Refrig until expiration date or RT for up to 28 days	RT for up to 28 days Do not refrigerate			Prime: 2 units Max: 60 units
	Humulin-N U-100 Vial		Refrig until expiration date or RT for up to 31 days	Refrig or RT for up to 31 days			-
	<a href="#">Humulin-N U-100 KwikPen</a>		Refrig until expiration date or RT for up to 14 days	RT for up to 14 days			Prime: 2 units Max: 60 units

All information obtained from Lexidrug or manufacturer package insert.  
Pen administration is illustrated at the bottom of each hyperlinked package insert.

RT: room temperature [68°F to 77°F (20°C to 25°C)]; Refrig: refrigeration [36°F - 46°F (2°C to 8°C)]. AC: before food; PC: after food.

	Label Name	Generic Name	Prior-Use Storage	In-Use Storage	When to Inject	Duration	Pen Priming & Max Injectable Dose
Long Acting	Lantus U-100 Vial	I. glargine	Refrig until expiration date or RT for up to 28 days	Refrig or RT for up to 28 days	Once daily at the same time each day	10.8 to >24 hours	-
	<a href="#">Lantus U-100 SoloStar</a>			RT for up to 28 days Do not refrigerate			Prime: 2 units Max: 80 units
	<a href="#">Toujeo/Insulin Glargine U-300</a> <a href="#">Solostar/Max Solostar</a>		Refrig until expiration date	RT for up to 56 days Do not refrigerate		>24 hours	Solostar: Prime: 3 units Max: 80 units  Max Solostar: Prime: 4 units Max: 160 units
Long Acting	<a href="#">Basaglar U-100 KwikPen</a>		Refrig until expiration date or RT for up to 28 days	RT for up to 28 days Do not refrigerate			Prime: 2 units Max: 80 units
	Tresiba/Insulin Degludec U-100 Vial	I. degludec	Refrig until expiration date or RT for up to 56 days	Refrig or RT for up to 56 days	Once daily (any time of day in adults; same time every day in pediatrics)	42 hours	-
	<a href="#">Tresiba/Insulin Degludec U-100 FlexTouch</a>						Prime: 2 units Max (U-100): 80 units Max (U-200): 160 units
	<a href="#">Tresiba/Insulin Degludec U-200 FlexTouch</a>						
	Semglee/Insulin Glargine-yfgn U-100 Vial	I. glargine-yfgn	Refrig until expiration or RT for up to 28 days	Refrig or RT for 28 days	Once daily at the same time each day	At least 24 hours	-
	<a href="#">Semglee/Insulin Glargine-yfgn U-100 Pen</a>			RT for up to 28 days Do not refrigerate			Prime: 2 units Max: 80 units
	<a href="#">Rezvoglar U-100 KwikPen</a>	I. glargine-aglr				> 24 hours	

All information obtained from Lexidrug or manufacturer package insert.  
Pen administration is illustrated at the bottom of each hyperlinked package insert.

RT: room temperature [68°F to 77°F (20°C to 25°C)]; Refrig: refrigeration [36°F - 46°F (2°C to 8°C)]. AC: before food; PC: after food.

	Label Name	Generic Name	Prior-Use Storage	In-Use Storage	When to Inject	Duration	Pen Priming & Max Injectable Dose
<b>70/30 Rapid acting mix</b>	Novolog Mix 70/30 Vial	I. aspart protamine/ I. aspart	Refrig until expiration date or RT for up to 28 days	Refrig or RT for up to 28 days	Within 15 mins AC (type 1) Within 15 mins AC/PC (type 2)	18 to 24 hours	-
	<a href="#">Novolog Mix 70/30 FlexPen</a>		Refrig until expiration date or RT for up to 14 days	RT for up to 14 days Do not refrigerate			Prime: 2 units Max: 60 units
<b>70/30 Short acting mix</b>	Novolin 70/30 Vial	I. NPH/ I. regular	Refrig until expiration date or RT for up to 42 days	RT for up to 42 days Do not refrigerate	Approximately 30 minutes AC (usually two divided doses)	Up to 24 hours	-
	<a href="#">Novolin 70/30 FlexPen</a>		Refrig until expiration date or RT for up to 28 days	RT for up to 28 days Do not refrigerate			Prime: 2 units Max: 60 units
	Humulin 70/30 Vial		Refrig until expiration date or RT for up to 31 days	Refrig or RT for up to 31 days	30 to 45 minutes AC (usually two divided doses)	18 to 24 hours	-
	<a href="#">Humulin 70/30 KwikPen</a>		Refrig until exp date or RT for up to 10 days	RT for up to 10 days Do not refrigerate			Prime: 2 units Max: 60 units
<b>75/25 Rapid acting mix</b>	Humalog Mix 75/25 Vial	I. lispro protamine/ I. lispro	Refrig until expiration date or RT for up to 28 days	Refrig or RT for up to 28 days	Within 15 mins AC (typically two divided doses)	14 to 24 hours	-
	<a href="#">Humalog Mix 75/25 KwikPen</a>		Refrig until expiration date or RT for up to 10 days	RT for up to 10 days Do not refrigerate			Prime: 2 units Max: 60 units
<b>50/50 Rapid acting mix</b>	<a href="#">Humalog Mix 50/50 KwikPen</a>						Prime: 2 units Max: 60 units
<b>U-500 Strength</b>	Humulin-R Vial	I. regular	Refrig until expiration date or RT for up to 40 days	Refrig or RT for up to 40 days	30 min AC	13 to 24 hours	-
	<a href="#">Humulin-R KwikPen</a>		Refrig until expiration date or RT for up to 28 days	RT for up to 28 days Do not refrigerate			Prime: 5 units Max: 300 units
<b>GLP-1/ Long-Acting Insulin Combo</b>	<a href="#">Soliqua</a>	I. glargine/ lixisenatide	Refrig until date expiration	RT for up to 28 days	Once daily within hour prior to first meal of the day	-	Prime: 2 units Max: 60 units
	<a href="#">Xultophy</a>	I. degludec/ liraglutide		RT or Refrig for up to 21 days	(Once daily/same time with or without food)	-	Max: 50 units Follow manufacturer instructions for priming

All information obtained from Lexidrug or manufacturer package insert.  
Pen administration is illustrated at the bottom of each hyperlinked package insert.

RT: room temperature [68°F to 77°F (20°C to 25°C)]; Refrig: refrigeration [36°F - 46°F (2°C to 8°C)]. AC: before food; PC: after food.